

ADDITIONS AND CORRECTIONS TO MY
REFERENCE LIST.

BY GREGORY M. MATHEWS.

299A. TADORNA TADORNOIDES WESTRALIS, subsp. n.

West Australia Sheld-drake.

Differs from *T. t. tadornoides* in being much less conspicuously marked and in its smaller size.

Type, South-west Australia.

Range, South-west Australia.

450A. APROSMICTUS ERYTHROPTERUS YORKI.

Little Crimson-winged Parrot.

Differs from *A. e. erythropterus* in its smaller size and from *A. e. coccinopterus* in being lighter coloured on the back.

Type, Cape York, No. 13,790.

Range, North Queensland.

555A. HALCYON PYRRHOPYGIUS UTINGI, subsp. n.

Allied Red-backed Kingfisher.

Differs from *H. p. obscurus* in having the wings more sky-blue, the head less green and the black nuchal stripe more pronounced.

Type, Cape York, No. 14,682.

Range, Cape York, North Queensland.

615. CHERAMŒCA LEUCOSTERNUM STONEL, subsp. n.

Black-and-White Swallow.

Differs from *C. c. leucosternum* in being much darker in the head and mantle.

Type, New South Wales, No. 1,549.

Range, New South Wales, Victoria, and South Australia.

663. *GERYGONE CULICIVORA BERNEYI*, subsp. n.
Queensland Fly-eater.

Differs from *G. c. exsul* in having a more pronounced buffish rump and a much smaller amount of white at the tip of the tail.

Type, Queensland, No. 14,514.

Range, Queensland.

- 870A. *CHTHONICOLA SAGITTATA QUEENSLANDICA* subsp. n.
Queensland Field-Wren.

Differs from *C. s. sagittata* in the almost entire absence of buff in the flanks, and in having a much smaller amount of white tipping to the tail-feathers.

Type, Queensland, No. 14,567.

Range, Queensland.

948. *MALURUS SPLENDENS RIORDANI*, subsp. n.
Northern Banded Wren.

Differs from *M. s. splendens* in its much deeper colour, and in having a very much slimmer bill.

Type, Yalgoo, Mid-Westralia, No. 5,250.

Range, Interior of Mid-Westralia.

- 1039A. *CRATICUS TORQUATUS COLEI*, subsp. n.
Mallee Butcher-Bird.

Differs from *C. t. olindus* in its lighter back and wing-coverts, and from *C. t. torquatus* in its smaller bill.

Type, Mallee, Victoria, No. 14,457.

Range, Mallee, Victoria.

1044. *FALCUNCULUS FRONTATUS LUMHOLTZI*, subsp. n.
Queensland Shrike-Tit.

Differs from *F. f. frontatus* in having a much shorter crest, more white in front of, and below the eye, and a much paler coloured tail with larger white tips.

Type, Queensland, No. 14,613.

Range, Queensland.

1325A. *ANTHUS AUSTRALIS QUEENSLANDICA*, subsp. n.
Northern Pipit.

Differs from *A. a. australis* in being darker above, with less white in the 2nd and 3rd tail-feathers. The shaft of the 2nd tail-feather is black, not brown.

Type, North Queensland, No. 14,623.

Range, Queensland.

1383A. *NEOCHMIA PHÆTON FITZROYI*, subsp. n.
Western Crimson Finch.

Differs from *N. p. phæton* in having the belly and the red on the throat lighter, in having a brown instead of a black head and in having a grey rump. From *N. p. iredalei* it differs in having the belly and the red on the throat darker and in having a much darker head.

Type, Fitzroy River, North-west Australia, No. 9,234.

Range, North-west Australia.

App. 3. *Ninox boobook royana*, subsp. n.
Norfolk Island Boobook.

Differs from *N. b. boobook* in its smaller size and the under surface covered with small spots.

Type, Norfolk Island.

Range, Norfolk Island.

385A. *NINOX CONNIVENS ADDENDA*, subsp. n.
South-western Winking Owl.

Differs from *N. c. connivens* in its larger size.

Type, South-west Australia.

Range, South-west Australia.